arXiv:0710.3508v1 [math.FA] 18 Oct 2007

WAVELET SETS WITHOUT GROUPS

MIHAELA DOBRESCU AND GESTUR ÓLAFSSON

Abstract. Most of the examples of wavelet sets are for dilation sets which are groups. We find a necessary and sufficient condition under which subspace wavelet sets exist for dilation sets of the form $\mathcal{AB}$, which are not necessarily groups. We explain the construction by a few examples.

Introduction

Wavelets and frames have become a widely studied tool in mathematics and applied science. One of the obvious questions is the construction of wavelets with given properties. The classical wavelet system on the line is given by a function $\psi \in L^2(\mathbb{R})$, such that the dyadic dilates and translates of ψ form an orthonormal basis for $L^2(\mathbb{R})$. Thus, $\{\psi_{j,n}\}_{j,n\in\mathbb{Z}}$ with $\psi_{j,n}(t) = 2^{j/2}\psi(2^j t + n)$, is an orthonormal basis for $L^2(\mathbb{R})$. There are several obvious generalizations: One can replace 2 by any integer N; one can allow several wavelet functions $\psi^1, \dots, \psi^L$; and one can consider an orthonormal basis for a closed subspace of $L^2(\mathbb{R})$. There have also been several publications of wavelets in higher dimensions, cf [1, 2, 3, 5, 10, 12, 13, 14, 21, 22, 23] to name few. One of the differences in higher dimensions is that we now have many more choices in the sets of dilations and translations. So, to fix the notation, let $\mathcal{D} \subseteq GL(n,\mathbb{R})$ and $\mathcal{T} \subseteq \mathbb{R}^n$ be countable sets. A $(\mathcal{D},\mathcal{T})$-*wavelet* is a square integrable function ψ with the property that the set of functions

$$\{|\det d|^{\frac{1}{2}}\varphi(dx+t) \mid d \in \mathcal{D}, t \in \mathcal{T}\} \tag{0.1}$$

forms an orthonormal basis for $L^2(\mathbb{R}^n)$. The set $\mathcal{D}$ is called the *dilation set* and the set $\mathcal{T}$ is called the *translation set*. If we replace $L^2(\mathbb{R}^n)$ in the above definition by

$$L^2_M(\mathbb{R}^n) = \{f \in L^2(\mathbb{R}^n) \mid \operatorname{supp}(\mathcal{F}(f)) \subseteq M\}$$

for some measurable subset $M \subseteq \mathbb{R}^n$, $|M| > 0$, we get a $(\mathcal{D},\mathcal{T})$-*s*ubspace wavelet. Here $\mathcal{F}$ stands for the Fourier transform

$$\mathcal{F}(f)(\lambda) = \int_{\mathbb{R}^n} f(t)e^{-2\pi i\langle t,\lambda\rangle}\, dt\,.$$

We will often write $\hat{f}$ for the Fourier transform of f.

The most natural starting point is to consider groups of dilations and full rank lattices as translation sets. The simplest examples would then be groups generated by one element $\mathcal{D} = \{a^k \mid k \in \mathbb{Z}\}$, see [24] and the reference therein. In [22, 23]

1991 *Mathematics Subject Classification.* 42C40,43A85.

Key words and phrases. Wavelet sets, spectral sets, tiling sets, subspace wavelets, wavelet transform.

M. Dobrescu was partially supported by DMS-0139783. The research of G. Ólafsson was supported by NSF grants DMS-0139783 and DMS-0402068.

more general sets of dilations were considered, and in general those dilations do not form a group. Even more general constructions can be found in [1].

In this article, we consider a special class of wavelets corresponding to *wavelet sets*. Those are wavelet functions ψ such that $\mathcal{F}(\psi) = \chi_\Omega$, for some measurable subset Ω of $\mathbb{R}^n$. The wavelet property is then closely related to geometric properties of the set Ω, in particular spectral and tiling properties of Ω. The study of wavelet sets then becomes an interplay between group theory, geometry, operator theory and analysis, cf [8, 6]. Our main results are existence theorems for such wavelets for some special dilation sets $\mathcal{D}$ which are *not necessarily* groups, see Theorem 2.1 and Theorem 2.3.

The article is organized as follows. In Section 1 we recall some well known facts on wavelet sets and their spectral and tiling properties. In Section 2 we give a construction of a special type of subspace wavelet set and we continue with two examples of such sets in Section 3. In Section 4 we discuss the general framework that has motivated the construction in this article. Those are the prehomogeneous vector spaces.

1. Wavelet sets

In this section we recall some basic facts about wavelet sets. In particular we discuss their relation to *spectral sets* and to *tilings*. A standard reference is the work by Y. Wang [26]. See also the discussion in [23] and the reference therein.

Set $e_\lambda(\xi) = e^{2\pi i \langle \lambda, \xi \rangle}$. If Ω is a Lebesgue measurable subset of $\mathbb{R}^n$, then

$$|\Omega| = \int \chi_\Omega(t_1, \ldots, t_n)\, dt_1 \ldots dt_n$$

denotes the measure of Ω with respect to the standard Lebesgue measure on $\mathbb{R}^n$.

Definition 1.1. A set $\Omega \subseteq \mathbb{R}^n$ with $0 < |\Omega| < \infty$ is a spectral set if there exists a set $\mathcal{T} \subseteq \mathbb{R}^n$ such that $\{e_\lambda \mid \lambda \in \mathcal{T}\}$ is an orthogonal basis for $L^2(\Omega)$. The set $\mathcal{T}$ is called a spectrum of Ω, and $(\Omega, \mathcal{T})$ is said to be a spectral pair.

Definition 1.2. A measurable tiling of a measure space (M, μ) is a countable collection of subsets $\{\Omega_j\}$ of M, such that

$$\mu(\Omega_i \cap \Omega_j) = 0\,,$$

for $i \neq j$, and

$$\mu(M \diagdown \bigcup_j \Omega_j) = 0\,.$$

Definition 1.3. Let $M \subseteq \mathbb{R}^n$ be a measurable set with $|M| > 0$. Let $\mathcal{D} \subseteq \mathrm{GL}(n, \mathbb{R})$ and $\mathcal{T} \subseteq \mathbb{R}^n$.

1) We call $\mathcal{D}$ a multiplicative tiling set of M if there exists a measurable set $\Omega \subseteq \mathbb{R}^n$, $|\Omega| > 0$, such that $\{d\Omega \mid d \in \mathcal{D}\}$ is a measurable tiling of M. The set Ω is called a multiplicative $\mathcal{D}$-tile for M or simply a *multiplicative tile*;
2) We call $\mathcal{T}$ an additive tiling set of $\mathbb{R}^n$ if there exists a measurable set $\Omega \subseteq \mathbb{R}^n$, $|\Omega| > 0$, such that $\{\Omega + t \mid t \in \mathcal{T}\}$ is a measurable tiling of $\mathbb{R}^n$. The set Ω is called an *additive $\mathcal{T}$-tile* or simply an *additive tile*;
3) A set Ω is called a *$(\mathcal{D}, \mathcal{T})$-tile* for M if Ω is a $\mathcal{D}$-multiplicative tile for M and a $\mathcal{T}$-additive tiling set for $\mathbb{R}^n$.

Remark 1.4. If $M = \mathbb{R}^n$ then we will simply speak of multiplicative tiling and $(\mathcal{D}, \mathcal{T})$-tile, without the reference to M.

Remark 1.5. Note, that we do not assume that $\mathcal{D}M \subseteq M$ or even

$$|(M \setminus \mathcal{D}M) \cup ((\mathcal{D}M) \setminus M)| = 0\,. \tag{1.1}$$

Neither do we assume that there exists a zero set Z, such that $\Omega \subseteq M \cup Z$. That will always be the case if $\mathrm{id} \in \mathcal{D}$. It should be noted, that we can always, without loss of generality, assume that $\mathrm{id} \in \mathcal{D}$ and $\Omega \subseteq M$. To see that, fix a $d \in \mathcal{D}$ and replace $\mathcal{D}$ by $\mathcal{D}d^{-1}$ and Ω by $d\Omega \subseteq M$ (up to a set of measure zero). Note, that if $\mathcal{D}$ is a group, then $\Omega \subseteq M$, and (1.1) follows from the tiling property (1).

The spectral property of a set is closely related to the tiling property, in particular if the spectrum is a lattice. This was first noticed by Fuglede in [11]. For a nonempty subset $\mathcal{T} \subset \mathbb{R}^n$, set

$$\mathcal{T}^* := \{t \in \mathbb{R}^n \mid \langle t, s\rangle \in \mathbb{Z},\ \text{for all } s \in \mathcal{T}\}\,.$$

If $\mathcal{T}$ is a lattice, then so is $\mathcal{T}^*$. In that case $\mathcal{T}^*$ is called *the dual lattice* of $\mathcal{T}$.

Theorem 1.6 (Fuglede [11])**.** *Assume that $\mathcal{T}$ is a lattice. Then Ω is a spectral set with spectrum $\mathcal{T}$ if and only if $\{\Omega + t \mid t \in \mathcal{T}^*\}$ is a measurable tiling of $\mathbb{R}^n$.*

This result and several examples led Fuglede to conjecture, cf. [11]:

Conjecture 1 (The Spectral-Tile Conjecture)**.** *A measurable set Ω, with positive and finite measure, is a spectral set if and only if it is an additive tile.*

Several people worked on this conjecture and derived important results and validated the conjecture for some special cases, see [15, 16, 17, 20, 26] and the references therein. However, in 2003, T. Tao [25] showed that the conjecture is false in dimension 5 and higher if the lattice hypothesis is dropped. The other direction was disproved by M. N. Kolountzakis and M. Matolcsi in [19, 18]. But even now, after the Spectral-Tiling conjecture has been proven to fail in higher dimensions, it is still interesting and important to understand better the connection between spectral properties and tiling in particular, because of the connection to wavelet sets.

Theorem 1.7 (Tao [25])**.** *Let $n \geq 5$ be an integer. Then there exists a compact set $\Omega \subset \mathbb{R}^n$ of positive measure such that $L^2(\Omega)$ admits an orthonormal basis of exponentials $\{e_\lambda \mid \lambda \in \mathcal{T}\}$ for some $\mathcal{T} \subset \mathbb{R}^n$, but such that Ω can not tile $\mathbb{R}^n$ by translation. In particular, Fuglede's conjecture is false in $\mathbb{R}^n$ for $n \geq 5$.*

We now discuss briefly some results by Dai, Larson, and Speegle [8]. For that assume that $\mathcal{G}$ and $\mathcal{T}$ are countable groups acting on a measure space (M, μ) by measurable automorphisms. We think of $\mathcal{G}$ as "dilation" group and $\mathcal{T}$ as translations. We write correspondingly the action of $\mathcal{G}$ by $m \mapsto gm$ and the action of $\mathcal{T}$ by $m \mapsto t + m$. Two sets E and F are said to be *$\mathcal{G}$-dilation congruent*, $E \sim_{\mathcal{G}} F$, if there exist measurable partitions $\{E_i\}$ and $\{F_i\}$ of E and F, respectively, such that $F_i = g_i E_i$ for some $g_i \in \mathcal{G}$. Similarly, two sets E and F are said to be *$\mathcal{T}$-translation congruent*, $E \sim_{\mathcal{T}} F$, if there exist measurable partitions $\{E_i\}$ and $\{F_i\}$ of E and F, respectively, such that $F_i = t_i + E_i$ for some $t_i \in \mathcal{T}$.

In 1996, the above three authors showed in [8], that wavelet sets exist for groups of dilations. They first introduce the notion of *abstract dilation-translation pair.*

Definition 1.8 (Dai-Larson-Speegle [9])**.** Let M be a metric space and $\mathcal{D}$ and $\mathcal{T}$ discrete groups of automorphisms of M. A pair $(\mathcal{D},\mathcal{T})$ is called an abstract dilation-translation pair if the following holds:

1) For each bounded set E and each open set F there exist $d \in \mathcal{D}$ and $t \in \mathcal{T}$ such that $t + E \subseteq d(F)$;
2) There is a fixed point θ for $\mathcal{D}$ such that for any neighborhood N of θ and for any bounded set E, there is an element $d \in \mathcal{D}$ such that $d(E) \subseteq N$.

In [8] the following is proved:

Theorem 1.9 (Dai-Larson-Speegle [8])**.** *Let M be a metric space and $(\mathcal{D},\mathcal{T})$ an abstract dilation-translation pair with θ a $\mathcal{D}$ fixed point as above. If E and F are bounded measurable sets in M such that E contains a neighborhood of θ and F has nonempty interior and is bounded away from θ, then there exists a measurable set $W \subseteq M$, $W \subseteq \cup_{d\in\mathcal{D}} d(F)$ which is $\mathcal{D}$-dilation congruent to F and $\mathcal{T}$-translation congruent to E.*

We now apply this abstract result to $\mathbb{R}^n$. If $d \in \mathrm{GL}(n,\mathbb{R})$, $\gamma \in \mathbb{R}^n$, and $\psi : \mathbb{R}^n \to \mathbb{C}$ is measurable, set

$$\psi_{d,\gamma}(x) = |\det d|^{1/2}\psi(dx+\gamma)\,. \tag{1.2}$$

Note that the Fourier transform of $\psi_{d,\gamma}$ is given by

$$\widehat{\psi_{d,\gamma}}(\lambda) = e^{2\pi i<\gamma,d^{-T}\lambda>}\widehat{\psi}(d^{-T}\lambda)\,. \tag{1.3}$$

Definition 1.10. Let $M \subseteq \mathbb{R}^n$ be measurable, $|M| > 0$, and $\mathcal{D} \subset \mathrm{GL}(n,\mathbb{R})$. Let $\mathcal{T} \subset \mathbb{R}^n$ be discrete. Then a measurable set $\Omega \subseteq M$ is called an M-subspace $(\mathcal{D},\mathcal{T})$-wavelet set if the set of functions $\{\psi_{d,\gamma}\}_{(d,\gamma)\in\mathcal{D}\times\mathcal{T}}$ is an orthogonal basis for $L^2_M(\mathbb{R}^n)$, where $\psi = \mathcal{F}^{-1}\chi_\Omega$.

Remark 1.11. Note again that we do not assume that $\Omega \subset M$, nor that $\mathcal{D}M = M$ up to set of measure zero, but this follows if $\mathrm{id} \in \mathcal{D}$. As in Remark 1.5 one can always assume this by replacing $\mathcal{D}$ by $d^{-1}\mathcal{D}$ and Ω by $d^T\Omega$ for a fixed $d \in \mathcal{D}$.

We get from Theorem 1.9:

Theorem 1.12 (Dai-Larson-Speegle [8])**.** *Let a be an expansive matrix, and let $M \subseteq \mathbb{R}^n$ be a measurable set of positive measure such that $a^TM = M$. Let $\mathcal{D} = \{a^k \mid k \in \mathbb{Z}\}$ and let $\mathcal{T}$ be a full rank lattice. Then there exists an M-subspace $(\mathcal{D},\mathcal{T})$-wavelet set.*

There are several generalizations of this Theorem. We refer to [23] for discussion and references. We will only mention two important results here:

Theorem 1.13 (Dai-Diao-Gu-Han [6])**.** *Let M be a measurable subset of $\mathbb{R}^n$, with positive measure satisfying $a^TM = M$, for some expansive matrix a, and let $\mathcal{T}$ be a full rank lattice. Then there exists a set $\Omega \subseteq M$ such that $\{\Omega + t \mid t \in \mathcal{T}\}$ is a measurable tiling of $\mathbb{R}^n$ and $\{(a^T)^k\Omega \mid k \in \mathbb{Z}\}$ is a measurable tiling of M. In particular Ω is an M-subspace $(\mathcal{D},\mathcal{T})$-wavelet set.*

Theorem 1.14 (Wang [26])**.** *Let $\mathcal{D} \subseteq GL(n,\mathbb{R})$ and $\mathcal{T} \subseteq \mathbb{R}^n$. Let $\Omega \subseteq \mathbb{R}^n$ be measurable, with positive and finite measure. If Ω is a measurable $\mathcal{D}^T$-tile and $(\Omega,\mathcal{T})$ is a spectral pair, then Ω is a $(\mathcal{D},\mathcal{T})$-wavelet set. Conversely, if Ω is a $(\mathcal{D},\mathcal{T})$-wavelet set and $0 \in \mathcal{T}$, then Ω is a measurable $\mathcal{D}^T$-tile and $(\Omega,\mathcal{T})$ is a spectral pair.*

Let us sketch some of the ideas of the proof to underline the connection between spectral properties, tiling, and wavelet sets. Let $\psi = \mathcal{F}^{-1}\chi_\Omega$. As the Fourier transform is an unitary isomorphism, it follows, that the set $\{\psi_{d,t} \mid d \in \mathcal{D}, t \in \mathcal{T}\}$ is an orthogonal basis for $L^2_M(\mathbb{R}^n)$ if and only if the set $\{\widehat{\psi_{d,t}} \mid d \in \mathcal{D}, t \in \mathcal{T}\}$ is an orthogonal basis for $L^2(M)$. Here, as before, we have set

$$\psi_{d,t}(x) = |\det d|^{1/2}\psi(dx + t)\,.$$

A simple calculation shows that

$$\widehat{\psi_{d,t}}(\lambda) = |\det d|^{-\frac{1}{2}} e^{2\pi i\langle d^{-1}t,\lambda\rangle}\chi_{d^T\Omega}(\lambda) = |\det d|^{-\frac{1}{2}} e^{2\pi i\langle t,d^{-T}\lambda\rangle}\chi_{d^T\Omega}(\lambda)\,.$$

The fact, that $d^T\Omega$ is a measurable tiling of M implies that

$$L^2(M) \simeq \bigoplus_{d\in\mathcal{D}} L^2(d^T\Omega)\,.$$

The orthogonal projection onto $L^2(d^T\Omega)$ is given by $f \mapsto f\chi_{d^T\Omega}$ and

$$f = \sum_{d\in\mathcal{D}} f\chi_{d^T\Omega}\,.$$

The spectral property implies, that $\{e_t\}_{t\in\mathcal{T}}$ is an orthogonal basis for $L^2(\Omega)$. As the linear map $f \mapsto |\det d|^{-1/2}f(d^{-T}\cdot)$ is a unitary isomorphism $L^2(\Omega) \simeq L^2(d^T\Omega)$ it follows, that the set of functions $\{|\det d|^{-1/2}e^{2\pi i\langle t,d^{-T}\cdot\rangle} = |\det d|^{-1/2}e_{d^{-1}t} \mid t \in \mathcal{T}\}$ is an orthogonal basis for $L^2(d^T\Omega)$. Putting those two things together, implies that $\{|\det d|^{-1/2}e_{d^{-1}t} \mid d \in \mathcal{D},\ t \in \mathcal{T}\}$ is an orthogonal basis for $L^2(M)$. Hence $\{\psi_{d,t} \mid (d,t) \in \mathcal{D}\times\mathcal{T}\}$ is an orthogonal basis for $L^2_M(\mathbb{R}^n)$.

We refer to [23] for further discussion and generalizations of Theorem 1.14.

2. Existence of subspace wavelet sets

In this section we discuss how to construct subspace wavelet sets using a kind of "induction" process, i.e., using well known facts discussed in the previous section on smaller dilation sets acting on a smaller frequency set and then extending those to our bigger dilation set and frequency set. We start with two simple, but important, observations. For $\mathcal{A},\mathcal{B} \subset \mathrm{GL}(n,\mathbb{R})$ we say that the product $\mathcal{AB} = \{ab \mid a \in \mathcal{A},\ b \in \mathcal{B}\}$ is *direct* if $a_1b_1 = a_2b_2$, $a_1,a_2 \in \mathcal{A}$, $b_1,b_2 \in \mathcal{B}$, implies that $a_1 = a_2$ and $b_1 = b_2$.

We state the following simple Lemma, but note, that we will be using the proof more than the actual statement. We remark, that the statements in the next few lemmas and theorems hold for more general settings, i.e., one could replace $\mathbb{R}^n$ by a measure space and $\mathrm{GL}(n,\mathbb{R})$ by a group of automorphisms of M.

Lemma 2.1. *Let $M \subseteq \mathbb{R}^n$ be measurable. Let $\mathcal{A},\mathcal{B} \subset \mathrm{GL}(n,\mathbb{R})$ be two non-empty sets, such that the product $\mathcal{AB}$ is direct Let $\mathcal{D} = \mathcal{AB} = \{ab \mid a \in \mathcal{A}\,,\ b \in \mathcal{B}\}$. Then there exists a $\mathcal{D}$-tile Ω for M if and only if there exists a measurable set $N \subseteq \mathbb{R}^n$, such that $\mathcal{A}N$ is a measurable tiling of M, and a $\mathcal{B}$-tile Ω for N.*

Proof. Assume that Ω is a $\mathcal{D}$-tile for M. Set

$$N := \mathcal{B}\Omega = \bigcup_{b\in\mathcal{B}} b\Omega\,.$$

Assume, that there are $b_1,b_2 \in \mathcal{B}$ such that $|b_1\Omega \cap b_2\Omega| > 0$. Then $|(ab_1\Omega) \cap (ab_2\Omega)| > 0$ for all $a \in \mathcal{A}$, which contradicts our assumption, that $\mathcal{D}\Omega$ is a measurable tiling of M. Hence $\mathcal{B}\Omega$ is a measurable tiling of N. We have, up to a set of measure zero:

$$\mathcal{A}N = \bigcup_{a\in\mathcal{A}} aN = \bigcup_{a\in\mathcal{A},\, b\in\mathcal{B}} ab\Omega = M\,.$$

Assume, that there are $a_1, a_2 \in \mathcal{A}$ such that $|a_1N \cap a_2N| > 0$. Then we can find $b_1, b_2 \in \mathcal{B}$ such that $|a_1b_1\Omega \cap a_2b_2\Omega| > 0$. As the product $\mathcal{AB}$ is direct, and $\mathcal{D}\Omega$ is a measurable tiling of M, it follows that $a_1b_1 = a_2b_2$ and hence $a_1 = a_2$. Thus $\mathcal{A}N$ is a measurable tiling of M.

For the other direction, assume that $\mathcal{A}N$ is a measurable tiling of M and that $\mathcal{B}\Omega$ is a measurable tiling of N. Then (up to sets of measure zero)

$$\bigcup_{d\in\mathcal{D}} d\Omega = \bigcup_{a\in\mathcal{A}} \bigcup_{b\in\mathcal{B}} ab\Omega = \bigcup_{a\in\mathcal{A}} a \bigcup_{b\in\mathcal{B}} b\Omega = \bigcup_{a\in\mathcal{A}} aN = M\,.$$

Assume that $|d_1\Omega \cap d_2\Omega| > 0$. Then there are unique $a_1, a_2 \in \mathcal{A}$, and $b_1, b_2 \in \mathcal{B}$ such that $d_1 = a_1b_1$ and $d_2 = a_2b_2$. Hence $|a_1N \cap a_2N| > 0$ which implies that $a_1 = a_2$, as $\mathcal{A}N$ is a measurable tiling for M. But then $|b_1\Omega \cap b_2\Omega| > 0$, which implies that $b_1 = b_2$. Hence $d_1 = d_2$. This shows, that $\mathcal{D}\Omega$ is a measurable tiling of M. □

Remark 2.2. We would like to remark at this point, that we do not assume that $\Omega \subseteq M$, nor that $N \subseteq M$. But this will in fact be the case in most applications because $\mathcal{D}$ will contain the identity matrix. Recall also from Remark 1.5 and Remark 1.11 that we can always assume that $\mathrm{id} \in \mathcal{D}$ and $\Omega \subseteq M$ up to set of measure zero. The same remarks hold for the following Theorems.

Theorem 2.3 (Construction of wavelet sets by steps, I)**.** *Let $\mathcal{M}, \mathcal{N} \subset \mathrm{GL}(n, \mathbb{R})$ be two non-empty subsets such that the product $\mathcal{MN}$ is direct. Let $\mathcal{L} = \mathcal{MN}$. Assume that $M \subseteq \mathbb{R}^n$ with $|M| > 0$, is measurable. Let $\mathcal{T} \subset \mathbb{R}^n$ be discrete. Then there exists a $(\mathcal{L}, \mathcal{T})$-wavelet set $\Omega \subset M$ for M if and only if there exists a $\mathcal{N}^T$-tiling set $N \subset M$ and a $(\mathcal{M}, \mathcal{T})$-wavelet set Ω_1 for N.*

Proof. Set $\mathcal{A} = \mathcal{N}^T$ and $\mathcal{B} = \mathcal{M}^T$. Then the conditions in Lemma 2.1 are satisfied.

Assume that $\Omega \subset M$ is a $(\mathcal{L}, \mathcal{T})$-wavelet set for M. As above we set

$$N := \mathcal{B}\Omega := \bigcup_{b\in\mathcal{M}} b^T\Omega\,.$$

Then, as above, we see that $\mathcal{A}N$ is a measurable tiling of M. As Ω is a spectral set, it follows from Theorem 1.14 that Ω is a $(\mathcal{M}, \mathcal{T})$-wavelet set for N.

Assume now that N is a $\mathcal{N}^T$-tiling for M, and that Ω_1 is a $(\mathcal{M}, \mathcal{T})$-wavelet set for N. Then, in particular Ω_1 is a $\mathcal{B}$-tile for N. As $\mathcal{A}N$ is a measurable tiling of M, it follows that $\mathcal{L}^T\Omega_1$ is a measurable tiling of M. As Ω_1 is a spectral set it follows from Theorem 1.14 that Ω_1 is a $(\mathcal{L}, \mathcal{T})$-wavelet set for M. □

Recall that if $\mathcal{D} \subseteq \mathrm{GL}(n, \mathbb{R})$, and $\mathcal{G} \subset \mathrm{GL}(n, \mathbb{R})$ is a group that acts on $\mathcal{D}$ from the right, then there exists a subset $\mathcal{D}_1 \subseteq \mathcal{D}$, such that $\mathcal{D} = \mathcal{D}_1\mathcal{G}$ and the product is direct. Note, that we do not assume that $\mathcal{G} \subset \mathcal{D}$.

Theorem 2.4 (Construction of wavelet sets by steps, II)**.** *Let $\mathcal{D} \subset \mathrm{GL}(n, \mathbb{R})$ and $M \subseteq \mathbb{R}^n$ measurable with $|M| > 0$. Let $\mathcal{T} \subset \mathbb{R}^n$ be discrete. Assume that $\mathcal{G} \subset \mathrm{GL}(n, \mathbb{R})$ is a group that acts on $\mathcal{D}$ from the right. Let $\mathcal{D}_1 \subseteq \mathcal{D}$ be such that $\mathcal{D} = \mathcal{D}_1\mathcal{G}$ as a direct product. Then there exists a $(\mathcal{D}, \mathcal{T})$-wavelet set Ω for M if only only if there exists a $\mathcal{G}^T$-tiling set N for M and a $(\mathcal{D}_1, \mathcal{T})$-wavelet set Ω_1 for N.*

Proof. This follows directly from Theorem 2.3 with $\mathcal{M} = \mathcal{D}_1$ and $\mathcal{N} = \mathcal{G}$. □

The question is then, how to obtain a wavelet set for the starting subset N. The following gives one way to do that.

Lemma 2.5 (Existence of subspace wavelet sets)**.** *Let $M \subseteq \mathbb{R}^n$ be a measurable set, $|M| > 0$. Let $a \in \mathrm{GL}(n, \mathbb{R})$ be an expansive matrix and $\emptyset \neq \mathcal{D} \subset \mathrm{GL}(n, \mathbb{R})$. Assume that $\mathcal{D}^T$ is a multiplicative tiling of M, $a\mathcal{D} = \mathcal{D}$ and $a^T M = M$. If $\mathcal{T}$ is a lattice, then there exists a measurable set $\Omega \subseteq M$ such that $\Omega + \mathcal{T}$ is a measurable tiling of $\mathbb{R}^n$ and $\mathcal{D}^T\Omega$ is a measurable tiling of M. In particular, Ω is a M-subspace $(\mathcal{D}, \mathcal{T})$-wavelet set.*

Proof. Let $b = a^T$ and $\mathcal{B} = \{b^k \mid k \in \mathbb{Z}\}$. Then $\mathcal{B}$ is an abelian group that acts on $\mathcal{D}^T$ from the right. Hence, there exists a set $\mathcal{A} \subset \mathcal{D}$ such that $\mathcal{D}^T = \mathcal{A}\mathcal{B}$ and the product is direct. Thus, the conditions in the previous Theorems are satisfied.

Let $E \subset M$ be such that $\mathcal{D}^T E$ is a measurable tiling of M. Set $N := \mathcal{B}E \subseteq M$. Then N is $\mathcal{B}$ invariant. By Theorem 1.12 there exists a $(\mathcal{B}, \mathcal{T})$-wavelet set Ω for N. Set

$$N := \bigcup_{k\in\mathbb{Z}} b^k\Omega\,.$$

Then, as before, we see that Ω is a $(\mathcal{D}, \mathcal{T})$-wavelet. □

3. Two simple examples

Before we discuss more general situations in the next section, let us explain our results by two simple examples.

Example 3.1. For $\theta \in \mathbb{R}$, let $R_\theta = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}$ denote the rotation in $\mathbb{R}^2$ by the angle θ. Let $a > 1$. For any integer $m \geq 2$ let

$$\mathcal{D}_{a,m} := \{a^n R^k_{2\pi/m} | n \in \mathbb{Z},\ k = 0, \ldots, m-1\}\,.$$

Note, that $R^k_{2\pi/m} = R_{2\pi k/m}$ and that $\mathcal{D}_{a,m}$ is a group. Let $\mathcal{T} = \mathbb{Z}^2$ and

$$\mathbb{R}^2_{2\pi/m} = \{r(\cos\psi, \sin\psi)^T \mid 0 \leq \psi \leq 2\pi/m\,,\ r > 0\}\,.$$

Then $\mathbb{R}^2_{2\pi/m}$ is a tiling set for the finite group $\{R^k_{2\pi/m} \mid k = 0, 1, \ldots, m-1\}$. As $a\cdot\mathrm{id}$ is expansive and $a\mathbb{R}^2_{2\pi/m} = \mathbb{R}^2_{2\pi/m}$, it follows from Theorem 1.12 that there exists a $(\mathcal{A} := \{a^k \cdot \mathrm{id} \mid k \in \mathbb{Z}\}, \mathcal{T})$-wavelet set Ω for $\mathbb{R}^2_{2\pi/m}$ and hence a $(\mathcal{D}_{a,m}, \mathcal{T})$-wavelet set for $\mathbb{R}^2$, see also Lemma 2.5. We show here how to construct such a wavelet set. Note that we only have to construct a $(\mathcal{A}, \mathcal{T})$-wavelet set for $\mathbb{R}^2_{2\pi/m}$. For that, let

$$E = [0,1] \times [0, \tan(2\pi/m)] \text{ if } m \neq 2, 4$$

$$E = [0,1]^2 \text{ if } m = 4$$

$$E = [-1,1] \times [0,1] \text{ if } m = 2$$

$$F = \{(x,y) \in \mathbb{R}^2_{2\pi/m} | 1 < x < a\}.$$

It is clear that $(\mathcal{D}_{a,m}, \mathcal{T})$ is an abstract dilation-translation pair with a fixed point 0. By Theorem 1 in [8], it follows that there exists a measurable set W such that W and E are *$\mathcal{T}$-translation congruent*, and W and F are *$\mathcal{D}$-dilation congruent*. On the other hand, F is a $\mathcal{D}_{a,m}$-multiplicative tile and $\{E, \mathcal{T}\}$ is a spectral pair. It



Figure 1. A $(\mathcal{D}_{2,4}, \mathcal{T})$ wavelet set.

follows that W is a $\mathcal{D}_{a,m}$-multiplicative tile and $\{W, \mathcal{T}\}$ is a spectral set. Thus, by Theorem 1.14 W is a $(\mathcal{D}_{a,m}, \mathcal{T})$ wavelet set.

.

Example 3.2. Let $A = \left(\begin{smallmatrix} 2 & 0 \\ 0 & 3 \end{smallmatrix}\right)$, $\mathcal{D} = \{R^k_{\pi/2} A^n \mid n \in \mathbb{Z}, k = 0, 1, 2, 3\}$ and $\mathcal{T} = \mathbb{Z}^2$. Note that $\mathcal{D}$ is not a group anylonger. Let $E = [0,1]^2$ and $F = ([0,2] \times [0,3]) \setminus [0,1]^2$. Then E is a $\mathbb{Z}^2$-additive tile and that F is a $\mathcal{D}$-multiplicative tile. Following the same procedure, we get a set W such that W and E are $\mathbb{Z}^2$-*translation congruent*, W and F are $\mathcal{D}$-*dilation congruent* and so, it follows that W is a $\mathcal{D}$-multiplicative tile and $\{W, \mathbb{Z}^2\}$ is a spectral set. Thus, W is a $(\mathcal{D}, \mathbb{Z}^2)$ wavelet set. The wavelet set W has the form

$$W = \bigcup_{i=1}^{2} \bigcup_{j=1}^{\infty} W_{i,j} \,.$$

The description of the set $W_{i,j}$ is as follows

$$\begin{aligned} W_{1,1} &= (E \setminus A^{-1}E) + (1,0) \\ W_{2,1} &= A^{-2}[(0,1) \times (1,3)] \end{aligned}$$

For $j \geq 2$, we have the following formulas

$$\begin{aligned} W_{1,j} &= [(A^{-j+1}E \setminus A^{-j}E) \setminus W_{2,j-1}] + (1,0) \\ W_{2,j} &= A^{-j-1}[W_{2,j-1} + (0,1)]. \end{aligned}$$

4. More general examples

The results in Section 2 give us tools to construct wavelet sets or subspace wavelet sets. In this section we discuss some general settings for our results. Those examples were discussed in [10, 22, 23] and the results therein were one of the main



Figure 2. The set W.

motivations for starting this work. We refer to the above articles for more information and details. The starting point is a subgroup $H \subseteq \mathrm{GL}(n, \mathbb{R})$ such that H^T has finitely many open orbits $U_1, \ldots, U_k$ in $\mathbb{R}^n$ such that

$$|\mathbb{R}^n \setminus (U_1 \cup \ldots \cup U_k)| = 0\,.$$

In this case, the pair $(H, \mathbb{R}^n)$ is called a *Prehomogeneous vector space*. Furthermore, we assume that

$$H^u = \{h \in H \mid h^T(u) = u\}$$

is compact for all $u \in U_j$, $j = 1, \ldots, k$. The reason for those assumptions is, that this implies that H is admissible in the sense of [21, 27], and we can define the continuous wavelet transform. In fact, there exists a function $\psi \in L^2(\mathbb{R}^n)$ such that

$$\int_H |\psi(h^T x)|^2 \, dh = 1$$

for almost all $x \in \mathbb{R}^n$. Here dh stands for a left-invariant Haar measure on H. Note, that we need the transpose of h here, because of the action in the frequency domain, cf (1.3).

Let $G := \mathbb{R}^n \times_s H$ be the semi-direct product of the groups $\mathbb{R}^n$ and H. This is a natural generalization of the $(ax + b)$-group in one dimension. Define $W_\psi : L^2(\mathbb{R}^n) \to L^2(\mathbb{R}^n \times_s H, \frac{dydh}{|\det h|})$ by

$$W_\psi(f)(y,h) = |\det h|^{-1/2} \int_{\mathbb{R}^n} f(x)\overline{\psi(h^{-1}(x-y))} \, dx \, . \tag{4.1}$$

That $W_\psi : L^2(\mathbb{R}^n) \to L^2(G)$ is an unitary isomorphism onto its image in $L^2(G)$ was shown in [10]. It is the *continuous wavelet transform.* The inverse of W_ψ is given by the weak integral:

$$f = \int_G W_\psi(f)(y,h)|\det h|^{-1/2}\psi(h^{-1}(\cdot - y) \, \frac{dydh}{|\det h|} \, . \tag{4.2}$$

We refer to [10] and Section 2 of [23] for detailed discussion and proofs. The natural question now arises to find a discrete version of the inversion formula (4.2). Note, that we can replace $L^2(\mathbb{R}^n)$ by the subspaces $L^2_{U_j}(\mathbb{R}^n)$ and define a continuous wavelet transform on each of those spaces. This is where the connection to the first part of this article comes into play.

Example 4.1. Let $n = 1$ and $H = \mathbb{R}^+$ acting on $\mathbb{R}$ by multiplication. Then G is just the $(ax + b)$-group. We have two open H-orbits $\mathbb{R}^+$ and $\mathbb{R}^- := -\mathbb{R}^+$. In each case the stabilizer of a point is trivial, i.e. just the one point subgroup $\{1\}$ and hence compact. In this case the space $L^2_{\mathbb{R}^+}(\mathbb{R})$ is the classical Hardy space of all functions in $L^2(\mathbb{R})$ that can be realized as L^2-boundary values of holomorphic functions on the upper half-plane $\mathbb{R} + i\mathbb{R}^+$. Similarly, the space $L^2_{\mathbb{R}^-}(\mathbb{R})$ is the space of all L^2-functions on the line, which can be written as L^2-boundary values of holomorphic functions on the lower half-plane $\mathbb{R} + i\mathbb{R}^-$. If we replace $\mathbb{R}^+$ by $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$, then there is only one open orbit $\mathbb{R}^*$.

Example 4.2. Let now $H = \mathbb{R}^+\mathrm{SO}(n)$. Then we have only one open orbit $\mathbb{R}^n \setminus \{0\}$. The stabilizer of a point in $\mathbb{R}^n \setminus \{0\}$ is naturally isomorphic to $\mathrm{SO}(n-1)$ and hence compact. A discrete version of the wavelet transform in this case would be a generalization of Example 3.1.

Example 4.3. Let $\Omega \subset \mathbb{R}^n$ be an open convex cone. Define

$$G(\Omega) := \{h \in \mathrm{GL}(n, \mathbb{R}) \mid h(\Omega) = \Omega\}$$

and

$$\Omega^* := \{w \in \mathbb{R}^n \mid \forall v \in \overline{\Omega} \setminus \{0\} \, : \, \langle w, v \rangle > 0\} \, .$$

The cone Ω is said to be homogeneous if $G(\Omega)$ acts transitively on Ω and self-dual if $\Omega^* = \Omega$. The cone Ω is symmetric if it is homogeneous and self-dual. Then there exists a solvable Lie group $H = AN \subset \mathrm{GL}(n, \mathbb{R})$, such that H acts freely and transitively on Ω. Here N is simply connected and nilpotent, the group A is simply connected and abelian, and A normalizes N. Note that $\mathbb{R}^+\mathrm{id} \subset A$ is a one dimensional subgroup containing an expansive matrix. Furthermore $\mathbb{R}^+\mathrm{id}$

commutes with all elements in H. This example is important in the study of Hardy spaces on $\mathbb{R}^n$ as well as Besov-spaces associated to symmetric cones, c.f., [4]

As in Section 3 of [23] we consider the case where the group H can be written as a semi direct product $H = ANR = NAR = RAN$ such that

1) R is compact, commutes with A, and normalizes N;
2) A is simply connected abelian and normalizing N;
3) the map
$$N \times A \times R \ni (n, a, r) \mapsto nar \in H$$
is a diffeomorphism;
4) There exists an expansive matrix $a \in A$ which is central in the group H;
5) there exists a co-compact discrete subgroup $\Gamma_N \subset N$.

In examples one of the groups can be trivial, and one could also use the group AR to fit the examples directly to the previous sections. Those assumptions includes all the examples above.

Assume now that M is one of the open H^T-orbits. For simplicity we assume that H^T acts freely on M, otherwise we will have to be a little more careful about the choice of Γ_R. Let $\mathbb{F}_N$ be a compact (or precompact, measurable) subset of N, such that $N = \mathbb{F}_N\Gamma_N$ is a measurable tiling of N. Let $\Gamma_A \subset A$ be a co-compact subgroup in A. Choose compact (or precompact measurable) subsets $\mathbb{F}_A \subset A$ such that $A = \mathbb{F}_A\Gamma_A$ is a measurable tiling. Finally let $\Gamma_R \subset R$ be a finite subgroup and $\mathbb{F}_R$ a compact (or precompact measurable) subset such that $\mathbb{F}_R\Gamma_R$ is a measurable tiling of R. Set $\mathbb{F}_H := \mathbb{F}_R\mathbb{F}_A\mathbb{F}_N$ and $\mathcal{D} := \Gamma_N\Gamma_A\Gamma_R$. Let $\mathbb{F} := \mathbb{F}_H^T m_0$, where $m_0 \in M$. Then $\mathcal{D}^T\mathbb{F}$ is a measurable tiling of H (c.f. the proof of Lemma 3.2 in [23]). Let a be the expansive matrix in (4) above. Then $a\mathcal{D} = \mathcal{D}$ and the assumptions in Lemma 2.5 are satisfied. It follows, that if $\mathcal{T}$ is any full rank lattice in $\mathbb{R}^n$, then there exists a $(\mathcal{D}, \mathcal{T})$-wavelet set for M. Using the corresponding wavelet function $\psi = |\Omega|^{-1/2}\mathcal{F}^{-1}\chi_\Omega$ allows us to write a discrete version of the inversion formula (4.2):

$$\begin{aligned} f &= \sum_{\gamma=(y,h)\in\mathcal{T}\times\mathcal{D}} W_\psi f(\gamma^{-1})\psi_{\gamma^{-1}} \\ &= \sum_{(y,h)\in\mathcal{T}\times\mathcal{D}} |\det h|^{1/2} W_\psi f(-h^{-1}y, h^{-1})\psi((\cdot) + y)\,. \end{aligned}$$

The factor $|\Omega|^{-1/2}$ is included to make ψ into an orthonormal wavelet function.

Note, that the assumption that the central element a as above exist is not satisfied for arbitrary groups H, even if H has finitely many open orbits, whose union is dense. As an example take the group

$$H = \left\{ \begin{pmatrix} a & t \\ 0 & 1/a \end{pmatrix} \mid a > 0,\, t \in \mathbb{R} \right\}$$

acting on $\mathbb{R}^2$. There are two open orbits, $y > 0$, and $y < 0$. Here A is the subgroup of diagonal matrices, and N is the subgroup of upper triangular matrices with 1s on the diagonal. In this case, the center of H is trivial. Note, that H is isomorphic to the $(ax + b)$-group, hence this is in fact an example of the $(ax + b)$-group acting on $\mathbb{R}^2$.

References

[1] A. Aldroubi, C. Cabrelli, and U. M. Molter. Wavelets on irregular grids with arbitrary dilation matrices and frame atoms for $L^2(\mathbb{R}^d)$. *Appl. Comput. Harmon. Anal.*, 17(2):119–140, 2004.

[2] P. Aniello, G. Cassinelli, E. De Vito, and A. Levrero. Wavelet transforms and discrete frames associated to semidirect products. *J. Math. Phys.*, 39(8):3965–3973, 1998.

[3] L. Baggett, A. Carey, W. Moran, and P. Ohring. General existence theorems for orthonormal wavelets, an abstract approach. *Publ. Res. Inst. Math. Sci.*, 31(1):95–111, 1995.

[4] D. Békollé, A. Bonami, G. Garrigós, and F. Ricci. Littlewood-Paley decompositions related to symmetric cones and Bergman projections in tube domains. *Proc. London Math. Soc. (3)*, 89(2):317–360, 2004.

[5] D. Bernier and K. F. Taylor. Wavelets from square-integrable representations. *SIAM J. Math. Anal.*, 27(2):594–608, 1996.

[6] X. Dai, Y. Diao, Q. Gu, and D. Han. The existence of subspace wavelet sets. *J. Comput. Appl. Math.*, 155(1):83–90, 2003. Approximation theory, wavelets and numerical analysis (Chattanooga, TN, 2001).

[7] X. Dai and D. R. Larson. Wandering vectors for unitary systems and orthogonal wavelets. *Mem. Amer. Math. Soc.*, 134(640):viii+68, 1998.

[8] X. Dai, D. R. Larson, and D. M. Speegle. Wavelet sets in $\mathbb{R}^n$. *J. Fourier Anal. Appl.*, 3(4):451–456, 1997.

[9] X. Dai, D. R. Larson, and D. M. Speegle. Wavelet sets in $\mathbb{R}^n$. II. In *Wavelets, multiwavelets, and their applications (San Diego, CA, 1997)*, volume 216 of *Contemp. Math.*, pages 15–40. Amer. Math. Soc., Providence, RI, 1998.

[10] R. Fabec and G. Ólafsson. The continuous wavelet transform and symmetric spaces. *Acta Appl. Math.*, 77(1):41–69, 2003.

[11] B. Fuglede. Commuting self-adjoint partial differential operators and a group theoretic problem. *J. Functional Analysis*, 16:101–121, 1974.

[12] H. Führ. Wavelet frames and admissibility in higher dimensions. *J. Math. Phys.*, 37(12):6353–6366, 1996.

[13] H. Führ. Continuous wavelet transforms with abelian dilation groups. *J. Math. Phys.*, 39(8):3974–3986, 1998.

[14] C. E. Heil and D. F. Walnut. Continuous and discrete wavelet transforms. *SIAM Rev.*, 31(4):628–666, 1989.

[15] A. Iosevich, N. H. Katz, and T. Tao. Convex bodies with a point of curvature do not have Fourier bases. *Amer. J. Math.*, 123(1):115–120, 2001.

[16] P. E. T. Jorgensen and S. Pedersen. Orthogonal harmonic analysis of fractal measures. *Electron. Res. Announc. Amer. Math. Soc.*, 4:35–42 (electronic), 1998.

[17] P. E. T. Jorgensen and S. Pedersen. Spectral pairs in Cartesian coordinates. *J. Fourier Anal. Appl.*, 5(4):285–302, 1999.

[18] M. N. Kolountzakis and M. Matolcsi. Complex Hadamard matrices and the spectral set conjecture. In *Proc. 7th International Conference on Harmonic Analysis and Partial Differential Equations (El Escorial), 2004*, to appear. (CA/11512).

[19] M. N. Kolountzakis and M. Matolcsi. Tiles with no spectra. *Forum Math.*, to appear. (CA/0406127).

[20] J. C. Lagarias and Y. Wang. Spectral sets and factorizations of finite abelian groups. *J. Funct. Anal.*, 145(1):73–98, 1997.

[21] R. S. Laugesen, N. Weaver, G. L. Weiss, and E.N. Wilson. A characterization of the higher dimensional groups associated with continuous wavelets. *J. Geom. Anal.*, 12(1):89–102, 2002.

[22] G. Ólafsson. Continuous action of Lie groups on $\mathbb{R}^n$ and frames. *Int. J. Wavelets Multiresolut. Inf. Process.*, 3(2):211–232, 2005.

[23] G. Ólafsson and D. Speegle. Wavelets, wavelet sets, and linear actions on $\mathbb{R}^n$. In *Wavelets, frames and operator theory*, volume 345 of *Contemp. Math.*, pages 253–281. Amer. Math. Soc., Providence, RI, 2004.

[24] D. Speegle. On the existence of wavelets for non-expansive dilation matrices. *Collect. Math.*, 54(2):163–179, 2003.

[25] T. Tao. Fuglede's conjecture is false in 5 and higher dimensions. *Math. Res. Lett.*, 11(2-3):251–258, 2004.

[26] Y. Wang. Wavelets, tiling, and spectral sets. *Duke Math. J.*, 114(1):43–57, 2002.

[27] G. Weiss and E. N. Wilson. The mathematical theory of wavelets. In *Twentieth century harmonic analysis—a celebration (Il Ciocco, 2000)*, volume 33 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 329–366. Kluwer Acad. Publ., Dordrecht, 2001.

E-mail address: `dobrescu@math.lsu.edu`

E-mail address: `olafsson@math.lsu.edu`

Both authors: Department of Mathematics, Louisiana State University, Baton Rouge, LA 70803, USA